


nutricomplex[®]

red 15-5-30

Water-soluble NPK 15-5-30
fertilizer with magnesium, sulfur and micronutrients



DESCRIPTION

Nutricomplex Red 15-5-30 is manufactured with high pure raw materials and enriched with chelated micro-nutrients. Nutricomplex Red 15-5-30 can be used in all fertigation systems and it's free of chloride, sodium and carbonates

MAIN BENEFITS

- Highly water soluble
- High pure raw materials
- Full and balanced crop nutrition
- Enriched with high quality Tradecorp EDTA chelated micro nutrients
- Safe to use in fertigation and foliar
- Free of chloride and sodium
- Acid pH & low EC
- Coloured formulas: easy to use
- Without any cake problem, improving handling and storage



nutricomplex® red 15-5-30



TECHNICAL SPECIFICATIONS

- Total nitrogen (N): 15% w/w
- Nitric nitrogen: 8.5% w/w
- Ammonium nitrogen: 3.5% w/w
- Ureic nitrogen: 3% w/w
- Phosphorus (P₂O₅) soluble in water: 5% w/w
- Potassium (K₂O) soluble in water: 30% w/w
- Magnesium (MgO) soluble in water: 2% w/w
- Sulphur (SO₃) soluble in water: 0.02% w/w
- Boron (B) soluble in water: 0.02% w/w
- Copper (Cu) soluble in water and chelated by EDTA: 0.01% w/w
- Iron (Fe) soluble in water and chelated by EDTA: 0.06% w/w
- Manganese (Mn) soluble in water and chelated by EDTA: 0.04% w/w
- Zinc (Zn) soluble in water and chelated by EDTA: 0.02% w/w

FOLIAR APPLICATION

Sugar beet	5 kg/ha/application	Apply at stage 4-6 leaves 2-3 times 10-14 days interval
Maize	5 kg/ha/application	Apply at stage 5-6 leaves 2 times with 7-10 days interval
Cereals	5 kg/ha/application	Apply in the autumn (frost protection), before and after flowering
Potato	5 kg/ha/application	Apply from 30 cm 2-3 times. 10-15 days interval
Oil seed rape	5 kg/ha/application	Apply in the autumn (frost protection) and the stage green buds
Tomato, paprika, cucumber	5 kg/ha/application	Apply during fruit setting and bulking 2-3 times within 10-14 days interval
Root vegetables	5 kg/ha/application	Apply during fruit bulking until 2-3 weeks before harvest
Cabbage, cauliflower	5 kg/ha/application	Apply during head/rose development. Maintain a regular interval of 2 weeks between 2 applications
Orchards	5 kg/ha/application	Apply during fruit bulking 3-5 times with 10-14 days of interval

DRIP IRRIGATION

Irrigation solution	0.5 to 2 g/L of irrigation water
Concentrated solution	15-20 kg/100 L of mother solution (100x concentrated)

DOSAGE AND METHOD OF USE