



tradecorp[®] AZ plus

Chemical mix combining
a complete mix of micro-elements and amino acids



DESCRIPTION

Chemical mix of essential micronutrients chelated with EDTA presented in the form of soluble micro-granules (WG) and enriched with active L- α -amino acids; indicated for preventing and correcting multiple micronutrient deficiencies

MAIN BENEFITS

- Our process of synthesis and chemical reaction ensures the maximum protection of the different micronutrients by the chelating agent
- Bio activator of enzymatic process of plants. To wake up and stimulate the plant at emergence from winter dormancy
- Very unique product containing active L- α -amino acids
- Total and instantaneous solubility without forming lumps or sediments
- Maximum quality and homogeneity since every micro-granule has the same composition and richness of chelated micronutrients
- Ideal formulation for fertigation, hydroponics and foliar application

TECHNICAL SPECIFICATIONS

- Iron (Fe) soluble in water and chelated by EDTA: 6% w/w
- Manganese (Mn) soluble in water and chelated by EDTA: 2.5% w/w
- Zinc (Zn) soluble in water and chelated by EDTA: 1.2% w/w
- Copper (Cu) soluble in water and chelated by EDTA: 0.6% w/w
- Boron (B) soluble in water: 0.5% w/w
- Molybdenum (Mo) soluble in water: 0.4% w/w
- Cobalt (Co) soluble in water and chelated by EDTA: 0.02% w/w
- Free amino acids: 2% w/w
- Chelating agent: EDTA
- Presentation: Soluble micro-granules (WG)
- Stable interval of pH of the chelated fraction: 3-10

tradecorp® AZ plus



SOIL APPLICATION

All crops	5-10 kg/ha	Apply before sowing, transplanting or beginning of plant growth
------------------	------------	---

HYDROPONIC CROPS

All crops	3-5 kg/1000 L solution	Use 1 L of stock solution/100 L irrigation water. Use the higher rate during crop development or periods of high temperature or during shorter and cooler days
------------------	------------------------	--

FERTIGATION

All crops	2-3 kg/ha	Apply every 7-10 days during the crop cycle, at periods of maximum needs
------------------	-----------	--

FOLIAR APPLICATION

General dosage	0.5-1.5 kg/ha	Apply when symptoms appear
Vegetables	0.25-1 kg/ha	3 applications. Apply at 10-15 days intervals, beginning when foliage is sufficient
Fruit trees, citrus, vines and olive trees	0.5-1 kg/ha	3 applications. Apply before flowering and after fruit setting
Herbaceous crops (cereals)	1 kg/ha	Apply during tillering, stem elongation and earing
Forage	1-1.5 kg/ha	Apply in each cutting
Potatoes and tuber vegetables	1 kg/ha	4 applications. Apply at 7-10 days intervals, beginning at 10 cm stage. Apply in a minimum of 500 L/ha of water
Ornamentals	0.25-0.5 kg/ha	Apply when symptoms appear

DOSAGE AND METHOD OF USE